

# Probability 2017

## 3

**Giovanni Pistone**

`www.giannidioresino.it`



DE CASTRO  
STATISTICS

**Collegio Carlo Alberto**

---

Mar 9, 2017

# Simple functions

Let  $(S, \mathcal{F})$  be a measurable space.

## Definition

A measurable real function,  $f: S \rightarrow \mathbb{R}$ ,  $f^{-1}: \mathcal{B} \rightarrow \mathcal{F}$ , is **simple** if it takes a finite number of values; equivalently, it is of the form  $f = \sum_{k=1}^m a_k \mathbf{1}_{A_k}$ ,  $a_k \in \mathbb{R}$ ,  $A_k \in \mathcal{F}$ ,  $k = 1, \dots, m$ ,  $m \in \mathbb{N}$ . The algebra with unity of all simple functions is denoted by  $\mathcal{S}$ ; the cone of all non-negative simple function is denoted by  $\mathcal{S}_+$ .

- Both  $\mathcal{S}$  and  $\mathcal{S}_+$  are closed for  $\vee$  and  $\wedge$ ;  $f = f^+ - f^-$ ,  $f \in \mathcal{S}$ .
- If  $f$  is measurable and non-negative, there exist an increasing sequence  $(f_n)_{n \in \mathbb{N}}$  in  $\mathcal{S}_+$  such that  $\lim_{n \rightarrow \infty} f_n(s) = f(s)$ ,  $s \in S$ .
- If  $f$  is measurable and bounded, there exist an sequence  $(f_n)_{n \in \mathbb{N}}$  in  $\mathcal{S}$  such that  $\lim_{n \rightarrow \infty} f_n = f$  uniformly.

## Integral of a non-negative function

Let  $(S, \mathcal{F}, \mu)$  be a measure space.

### Definition

- If  $f \in \mathcal{S}$ ,  $f = \sum_{k=1}^m a_k \mathbf{1}_{A_k}$ , we define its **integral** to be

$$\int f \, d\mu = \sum_{k=1}^m a_k \mu(A_k) \quad \text{where } 0 \cdot \infty = \infty \cdot 0 = 0$$

- If  $f: S \rightarrow [0, +\infty]$  is measurable, namely  $f \in \mathcal{L}_+$ , we define its **integral** to be

$$\int f \, d\mu = \sup \left\{ \int h \, d\mu \mid h \in \mathcal{S}_+, h \leq f \right\}$$

- The integral is linear and monotone on  $\mathcal{S}^1 = \{f \in \mathcal{S} \mid \int f^+ \, d\mu, \int f^- \, d\mu \leq \infty\}$ .

- The integral is convex and monotone on  $\mathcal{L}_+$ .




- If  $f \in \mathcal{L}_+$  and  $\int f \, d\mu = 0$ , then  $\mu\{f > 0\} = 0$ .

# Monotone-Convergence Theorem

## Theorem (MON)

let  $(f_n)_{n \in \mathbb{N}}$  be a non-decreasing sequence in  $\mathcal{L}_+$ . Then the pointwise limit  $f = \lim_{n \rightarrow \infty} f_n$  belongs to  $\mathcal{L}_+$  and  $\lim_{n \rightarrow \infty} \int f_n d\mu = \int f d\mu$ .

- A sequence of simple functions converging to  $f$  is always available.

- If  $\alpha, \beta \in \mathbb{R}_{>0}$ ,  $f, g \in \mathcal{L}_+$ , then :

$$\int (\alpha f + \beta g) d\mu = \int \alpha f d\mu + \int \beta g d\mu$$

- Exercise: If  $\mu(S) = 1$ , then  $\int f d\mu = \int_0^\infty \mu\{f > u\} du$ .

Proof of MON: Appendix A5 of D. Williams. *Probability with martingales*. Cambridge Mathematical Textbooks. Cambridge University Press, Cambridge, 1991; Bertrand Lods *Lecture Notes*.

# Fatou Lemmas

## Theorem (FATOU)

Let  $(f_n)_{n \in \mathbb{N}}$  be a sequence in  $\mathcal{L}_+$ .

1.  $\int (\liminf_{n \rightarrow \infty} f_n) d\mu \leq \liminf_{n \rightarrow \infty} \int f_n d\mu.$
2. If, moreover,  $f_n \leq g$ ,  $n \in \mathbb{N}$ , and  $\int g d\mu < \infty$ , then  $\int (\limsup_{n \rightarrow \infty} f_n) d\mu \geq \limsup_{n \rightarrow \infty} \int f_n d\mu.$

- Exercise: Prove 1. by observing that  $\limsup$  is the limit of an increasing sequence.
- Exercise: If  $(f_n)_{n \in \mathbb{N}}$  is a decreasing sequence in  $\mathcal{L}_+$  and  $\int f_1 d\mu < \infty$ , then  $\int (\lim_{n \rightarrow \infty} f_n) d\mu = \lim_{n \rightarrow \infty} \int f_n d\mu.$
- Exercise: If  $(f_n)_{n \in \mathbb{N}}$  is a sequence in  $\mathcal{L}_+$  and  $f_n \leq g$ ,  $n \in \mathbb{N}$ ,  $\int g d\mu < \infty$ , then  $\int (\lim_{n \rightarrow \infty} f_n) d\mu = \lim_{n \rightarrow \infty} \int f_n d\mu.$

# Integrability

## Definition

- Let  $\mathcal{L}^1$  be the vector space of measurable real functions such that

$$\int |f| d\mu = \int f^+ d\mu + \int f^- d\mu < \infty$$

- Define the integral to be the linear mapping

$$\mathcal{L}^1 \ni f \mapsto \int f d\mu = \int f^+ d\mu - \int f^- d\mu \in \mathbb{R}$$

- Jensen inequality: Let  $\Phi: \mathbb{R} \rightarrow \mathbb{R}$  be convex. Then  $\Phi\left(\int f d\mu\right) \leq \int \Phi \circ f d\mu$ .
- Assignment: Revise  $L^1$ -convergence and Dominated Convergence Theorem in Ch 5 of D. Williams. *Probability with martingales*. Cambridge Mathematical Textbooks. Cambridge University Press, Cambridge, 1991 or Bertrand Lods *Lecture Notes*.

## Change of variable formula

Let be given a measure space  $(S, \mathcal{F}, \mu)$ , a measurable space  $(\mathbb{X}, \mathcal{G})$  and a measurable mapping  $\phi: S \rightarrow \mathbb{X}$ ,  $\phi^{-1}: \mathcal{G} \rightarrow \mathcal{F}$ . Let  $\phi_{\#}\mu = \mu \circ \phi^{-1}$  be the push-forward measure.

- If  $h \in \mathcal{S}(\mathbb{X}, \mathcal{G})$  i.e.  $h = \sum_{k=1}^n b_k \mathbf{1}_{B_k}$ ,  $b_k \in \mathbb{R}$ ,  $B_k \in \mathcal{G}$ ,  $k = 1, \dots, n$ .  
Then

$$\begin{aligned} \int h \, d\phi_{\#}\mu &= \sum_{k=1}^n b_k \phi_{\#}\mu(B_k) = \sum_{k=1}^n b_k \mu[\phi^{-1}(B_k)] = \\ &= \sum_{k=1}^n b_k \int \mathbf{1}_{B_k} \circ \phi \, d\mu = \int h \circ \phi \, d\mu \end{aligned}$$

- If  $f \in \mathcal{L}_+$ , then MON implies  $\int h \, d\phi_{\#}\mu = \int h \circ \phi \, d\mu$
- If  $f: \mathbb{X} \rightarrow \mathbb{R}$  is measurable and  $f \circ \phi \in \mathcal{L}^1(S, \mathcal{F}, \mu)$  then  $f \in \mathcal{L}^1(\mathbb{X}, \mathcal{G}, \phi_{\#}\mu)$  and  $\int h \, d\phi_{\#}\mu = \int h \circ \phi \, d\mu$ .

## Densities

Let be given a measure space  $(S, \mathcal{F}, \mu)$  and a measurable non-negative mapping  $p: S \rightarrow \mathbb{R}$  such that  $\int p \, d\mu < \infty$ .

The set function

$$p \cdot \mu: \mathcal{F} \rightarrow \mathbb{R}_{>0}, \quad A \mapsto \int \mathbf{1}_A p \, d\mu$$

is a bounded measure. In fact:  $p \cdot \mu(\emptyset) = \int 0 \, d\mu = 0$ ; given a sequence  $(A_n)_{n \in \mathbb{N}}$  of disjoint events, then MON implies

$$\begin{aligned} p \cdot \mu(\cup_{n \in \mathbb{N}} A_n) &= \int \mathbf{1}_{\cup_{n \in \mathbb{N}} A_n} p \, d\mu = \int \sum_{n \in \mathbb{N}} \mathbf{1}_{A_n} p \, d\mu = \\ &= \sum_{n \in \mathbb{N}} \int \mathbf{1}_{A_n} p \, d\mu = \sum_{n \in \mathbb{N}} p \cdot \mu(A_n) \end{aligned}$$

Exercise: If  $f: S \rightarrow \mathbb{R}$  is measurable and  $fp \in \mathcal{L}^1(S, \mathcal{F}, \mu)$ , then  $f \in \mathcal{L}^1(S, \mathcal{F}, p \cdot \mu)$  and  $\int f \, d(p \cdot \mu) = \int fp \, d\mu$ . [Hint: try first simple functions, then use MON]



# Product measure I

## Definition

Given measure spaces  $(S_i, \mathcal{F}_i, \mu_i)$ ,  $i = 1, \dots, n$ ,  $n = 2, 3, \dots$ , the product measure space is

$$(S, \mathcal{F}, \mu) = \otimes_{i=1}^n (S_i, \mathcal{F}_i, \mu_i) = (\times_{i=1}^n S_i, \otimes_{i=1}^n \mathcal{F}_i, \otimes_{i=1}^n \mu_i),$$

where

$$\mathcal{F} = \otimes_{i=1}^n \mathcal{F}_i = \sigma \{ \times_{i=1}^n A_i \mid A_i \in \mathcal{F}_i, i = 1, \dots, n \}$$

and  $\mu = \otimes_{i=1}^n \mu_i$  is the unique measure on  $(\times_{i=1}^n S_i, \otimes_{i=1}^n \mathcal{F}_i)$  such that

$$\mu(\times_{i=1}^n A_i) = \prod_{i=1}^n \mu_i(A_i), \quad A_i \in \mathcal{F}_i, i = 1, \dots, n.$$

- Let  $X_i: S \mapsto S_i$ ,  $i = 1, \dots, n$ , be the projections. Then  $\otimes_{i=1}^n \mathcal{F}_i = \sigma \{ X_i \mid i = 1, \dots, n \}$ .
- Examples: Counting measure on  $\mathbb{N}^2$ , Lebesgue measure on  $\mathbb{R}^2$ , the finite Bernoulli scheme.
- Product measure of probability measures is a probability measure.

## Product measure II

Recall all measures are  $\sigma$ -finite. Assume  $n = 2$ .

### Sections

If  $C \in \mathcal{F} = \mathcal{F}_1 \otimes \mathcal{F}_2$ , then for each  $x_1 \in S_1$  the set  $\{x_2 \in S_2 \mid (x_1, x_2) \in C\}$  belongs to  $\mathcal{F}_2$ .

### Proof.

Let  $\mathcal{O}$  be the family of all subsets of  $S$  for which the proposition is true.  $\mathcal{O}$  is a  $\sigma$ -algebra that contains all the measurable rectangles, hence  $\mathcal{F} \subset \mathcal{O}$ . □

### Partial integration

The mapping  $S_1: x_1 \mapsto \mu_2 \{x_2 \in S_2 \mid (x_1, x_2) \in C\}$  is non-negative and  $\mathcal{F}_1$ -measurable.

### Proof.

If  $C \in \mathcal{F}$  then the function is well defined. The set of all  $C \in \mathcal{F}$  such that the function is measurable contains measurable rectangles, is a  $\pi$ -system, and is a  $d$ -system. □

## Product measure III

### Product measure: existence

The set function  $\mu: \mathcal{F} \ni C \mapsto \int \mu_2 \{x_2 \in S_2 | (x_1, x_2) \in C\} \mu_1(dx_1)$  is a measure such that  $\mu(A_1 \times A_2) = \mu_1(A_1)\mu_2(A_2)$  on measurable rectangles. Hence,  $\mu = \mu_1 \otimes \mu_2$ .

### Proof.

The integral exists because the integrand is non-negative.  $\mu(\emptyset) = 0$ ; if  $(C_n)_{n \in \mathbb{N}}$  is a sequence in  $\mathcal{F}$  of disjoint events, then for all  $x_1 \in S_1$  we have

$$\begin{aligned} \mu_2 \{x_2 \in S_2 | (x_1, x_2) \in \cup_{n \in \mathbb{N}} C_n\} &= \mu_2 (\cup_{n \in \mathbb{N}} \{x_2 \in S_2 | (x_1, x_2) \in C_n\}) = \\ &= \sum_{n \in \mathbb{N}} \mu_2 \{x_2 \in S_2 | (x_1, x_2) \in C_n\} \end{aligned}$$

MON implies

$$\begin{aligned} \mu (\cup_{n \in \mathbb{N}} C_n) &= \int \sum_{n \in \mathbb{N}} \mu_2 \{x_2 \in S_2 | (x_1, x_2) \in C_n\} \mu_1(dx_1) = \\ &= \sum_{n \in \mathbb{N}} \int \mu_2 \{x_2 \in S_2 | (x_1, x_2) \in C_n\} \mu_1(dx_1) = \sum_{n \in \mathbb{N}} \mu(C_n) \end{aligned}$$

## Product measure IV

- Consider  $n = 3$ . The product measure space

$$\otimes_{i=1}^3 (S_i, \mathcal{F}_i, \mu_i) = (\times_{i=1}^n S_i, \otimes_{i=1}^n \mathcal{F}_i, \otimes_{i=1}^n \mu_i)$$

is identified with

$$(S_1 \times S_2, \mathcal{F}_1 \otimes \mathcal{F}_2, \mu_1 \otimes \mu_2) \otimes (S_3, \mathcal{F}_3, (\mu_1 \otimes \mu_2) \otimes \mu_3)$$

One has to check that

$$(\mathcal{F}_1 \otimes \mathcal{F}_2) \otimes \mathcal{F}_3 = \mathcal{F}_1 \otimes \mathcal{F}_2 \otimes \mathcal{F}_3$$

- The  $n = \infty$  case requires Carathéodory. See the Bernoulli scheme example.

# Fubini theorem I

## Section

Let  $f: S_1 \times S_2 \rightarrow \mathbb{R}$  be  $\mathcal{F}_1 \otimes \mathcal{F}_2$  measurable. For all  $x_1 \in S_1$  the function  $f_{x_1}: x_2 \mapsto f(x_1, x_2)$  is  $\mathcal{F}_2$ -measurable.

## Proof.

For each  $y \in \mathbb{R}$ , consider the level set

$C = \{(x_1, x_2) \mid f(x_1, x_2) \leq y\} \in \mathcal{F}_1 \otimes \mathcal{F}_2$ . The set  $\{x_2 \mid f_{x_1}(x_2) \leq y\}$  is the  $x_1$ -section of  $C$ . □

## Theorem (Non-negative integrand)

Let  $f: S_1 \times S_2 \rightarrow \mathbb{R}$  be  $\mathcal{F}_1 \otimes \mathcal{F}_2$ -measurable and non-negative. Then the mapping  $S_1 \ni x_1 \mapsto \int f(x_1, x_2) \mu_2(dx_2)$  is  $\mathcal{F}_1$ -measurable and

$$\int f \, d\mu_1 \otimes \mu_2 = \int \left( \int f(x_1, x_2) \mu_2(dx_2) \right) \mu_1(dx_1)$$

## Fubini theorem II

### Theorem (Integrable integrand)

Let  $f: S_1 \times S_2 \rightarrow \mathbb{R}$  be  $\mu_1 \otimes \mu_2$ -integrable. Then the mapping  $S_1 \ni x_1 \mapsto \int f(x_1, x_2) \mu_2(dx_2)$  is  $\mu_1$ -integrable and

$$\int f d\mu_1 \otimes \mu_2 = \int \left( \int f(x_1, x_2) \mu_2(dx_2) \right) \mu_1(dx_1)$$

### Proof: Non-negative integrand.

Choose an increasing sequence of simple non-negative functions converging to  $f$  and use MON. □

### Proof: Integrable integrand.

Decompose  $f = f^+ - f^-$  and use the previous form of the theorem. □

# Independence

## Definition

Let  $(\Omega, \mathcal{F}, \mu)$  be a probability space.

1. The sub- $\sigma$ -algebras  $\mathcal{F}_1, \dots, \mathcal{F}_n$  are independent if  $A_i \in \mathcal{F}_i$ ,  $i = 1, \dots, n$ , implies  $\mu(A_1 \times \dots \times A_n) = \mu(A_1) \cdots \mu(A_n)$ .
2. The random variables  $X_i: \Omega \rightarrow S_i$ ,  $X_i^{-1}: \mathcal{G}_i \rightarrow \mathcal{F}$ ,  $i = 1, \dots, n$ , are independent, if

$$(X_1, \dots, X_n)_{\#}\mu = (X_1)_{\#}\mu \otimes \cdots \otimes (X_n)_{\#}\mu$$

- If  $\mathcal{F}_i = \sigma(X_i)$ , the 1. and 2. are equivalent.